STANDING ORDER CERTIFICATION FORM

Practitioner Name: Dr. Muntu Davis
Practitioner Address: **1000 Broadway, Suite 500, Oakland, CA 94607**
Practitioner Phone: **510-567-8010**
Date: **5/26/17**

Shipment Information

Recipient Name: **Amy Taxdahl, BSN, RN**
Recipient Title: **School Nurse**
Recipient Address: **San Leandro Unified School District, 2255 Bancroft Ave, San Leandro, CA 94577**

By signing this Certification form, I certify that, under applicable state law, I may (1) purchase and possess quantities of EpiPen® (epinephrine injection, USP) or EpiPen Jr® (epinephrine injection, USP) Auto-Injectors (the "Product"), (2) prescribe, furnish and administer the Product to patients, (3) issue standing orders or otherwise delegate to appropriate persons the authority to prescribe, furnish or administer the Product to a student at a school or involved in a school-related activity, and (4) ship, or direct shipment of, quantities of the Product to a school for prescribing, furnishing or administering to a student at the school or involved in a school-related activity.

Practitioner Signature: _____ MD, MPH
Date: August 11, 2017

Indications
EpiPen® (epinephrine injection, USP) 0.3 mg and EpiPen Jr® (epinephrine injection, USP) 0.15 mg Auto-Injectors are indicated in the emergency treatment of type 1 allergic reactions, including anaphylaxis, to allergens, idiopathic and exercise-induced anaphylaxis, and in patients with a history or increased risk of anaphylactic reactions. Selection of the appropriate dosage strength is determined according to body weight.

Important Safety Information
EpiPen® and EpiPen Jr® Auto-Injectors are intended for immediate administration as emergency supportive therapy only and are not intended as a substitute for immediate medical or hospital care. **In conjunction with the administration of epinephrine, the patient should seek immediate medical or hospital care**. More than two sequential doses of epinephrine should only be administered under direct medical supervision.

EpiPen® and EpiPen Jr® should **only** be injected into the anterolateral aspect of the thigh. **Do not inject intravenously, into buttock, or into digits, hands, or feet.** Instruct caregivers to hold the leg of young children firmly in place and limit movement prior to and during injection to minimize risk of injection-related injury.

Rare cases of serious skin and soft tissue infections have been reported following epinephrine injection. Advise patients to seek medical care if they develop symptoms of infection such as persistent redness, warmth, swelling, or tenderness at the injection site.

Epinephrine should be used with caution in patients with heart disease, and in patients who are on drugs that may sensitize the heart to arrhythmias, because it may precipitate or aggravate angina pectoris and produce ventricular

arrhythmias. Arrhythmias, including fatal ventricular fibrillation, have been reported, particularly in patients with underlying cardiac disease or taking cardiac glycosides, diuretics, or anti-arrhythmics.

Patients with certain medical conditions or who take certain medications for allergies, depression, thyroid disorders, diabetes, and hypertension, may be at greater risk for adverse reactions. Common adverse reactions to epinephrine include anxiety, apprehensiveness, restlessness, tremor, weakness, dizziness, sweating, palpitations, pallor, nausea and vomiting, headache, and/or respiratory difficulties.

Please see the full Prescribing Information and Patient Information.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

For additional information please contact us at 800-395-3376.

EpiPen®, EpiPen Jr® and *EpiPen4Schools*® are registered trademarks owned by the Mylan companies.

© 2016 Mylan Specialty L.P. All rights reserved.